

Lorentz-Covariant Green-Schwarz Superstring Amplitudes

Nathan Berkovits

Math Dept., King's College, Strand, London, WC2R 2LS, United Kingdom

e-mail: udah101@oak.cc.kcl.ac.uk

KCL-TH-92-6

November 1992

Abstract

In a recent paper, the BRST formalism for the gauge-fixed $N=2$ twistor-string was used to calculate Green-Schwarz supersring scattering amplitudes with an arbitrary number of loops and external massless states. Although the gauge-fixing procedure preserved the worldsheet $N=2$ superconformal invariance of the twistor-string, it broke the target-space $SO(9,1)$ super-Poincaré invariance down to an $SU(4) \times U(1)$ subgroup.

In this paper, generators for the $SO(9,1)$ super-Poincaré transformations, as well as manifestly covariant vertex operators, are explicitly constructed out of the gauge-fixed matter fields. The earlier calculated amplitudes are then expressed in manifestly Lorentz-covariant notation.

I. Introduction

There are various methods for calculating superstring amplitudes, each having advantages and disadvantages. The most common method is to start with the covariant Neveu-Schwarz-Ramond action in which all gauge invariances have been fixed except for the worldsheet $N=1$ superconformal invariance. Since this action contains only free fields, it is straightforward to calculate superstring scattering amplitudes by defining vertex operators for the physical states, evaluating correlation functions of these vertex operators on $N=1$ super-Riemann surfaces, integrating over the super-moduli of the surfaces, and finally summing over all possible spin structures.^{1,2} These scattering amplitudes can be proven to be unitary by showing agreement with amplitudes obtained using the light-cone NSR method.³

One disadvantage of this approach is that until the final step of summing over spin structures, the scattering amplitude is not spacetime supersymmetric, and therefore contains divergences coming from the dilaton tadpole diagram. In order to regularize these divergences, a cutoff in the moduli space must be introduced which can only be removed after summing over spin structures. Because the integrand of the scattering amplitude changes by a total derivative for different choices of anti-commuting moduli for the surface,² this cutoff produces a boundary term that depends on the choice of the anti-commuting moduli.⁴ Although an unambiguous choice for the anti-commuting moduli can be determined from unitary requirements,⁵ the need to make such a choice complicates the analysis of the multiloop scattering amplitudes.⁶ A second disadvantage of the NSR approach is that the fermionic vertex operator is a function of ghost fields, as well as matter fields.¹ This ghost dependence makes it awkward to formulate the NSR string in a fermionic background, preventing a straightforward derivation of the ten-dimensional supergravity equations of motion.

Another method for calculating superstring scattering amplitudes is to start with the light-cone Green-Schwarz action in which all gauge invariances including conformal invariance have been fixed.^{7,8,9} This manifestly unitary method differs in three important features from the covariant Neveu-Schwarz-Ramond method. Firstly, there is no sum over spin structures since spacetime supersymmetry is manifest. Secondly, the action is defined on ordinary Riemann surfaces, rather than on $N=1$ super-Riemann surfaces. And thirdly, non-trivial operators need to be inserted at the interaction points of the surface in order to preserve Lorentz invariance (these interaction points are located at the zeros of $\partial_z \rho$, where $\partial_z \rho$ is the unique meromorphic function with poles of residue P_r^{9+0} at $z = z_r$ and purely imaginary periods when integrated around the $2g$ non-trivial cycles).

Although the first two features of the light-cone Green-Schwarz method remove the problems of the covariant NSR method, the third feature introduces new problems. Since correlation functions now depend on the locations of the interaction points, as well as the locations of the vertex operators, the resulting expressions for the scattering amplitudes are complicated functions of the moduli of the surface. Furthermore, potential divergences when two or more interaction points coalesce force the introduction of contact

terms,^{9,10,11} whose contribution to multiloop Green-Schwarz amplitudes⁹ has not yet been included in any manifestly Lorentz-covariant expressions.

It should be noted that the semi-light-cone Green-Schwarz method,¹² in which all gauge invariances are fixed except for conformal invariance, has exactly the same advantages and disadvantages as the light-cone Green-Schwarz method.¹³ The only difference is that in order to preserve Lorentz invariance, the non-trivial operators need to be inserted at the zeros of the expectation value of $\partial_z x^{9+0}$, since if one fixes the remaining conformal invariance to get to light-cone gauge, $\partial_z x^{9+0}$ is replaced by $\partial_z \rho$.

Recently, a new method for calculating superstring amplitudes has been developed in which one starts from an N=2 twistor-string version of the Green-Schwarz action where all gauge invariances have been fixed except for worldsheet N=2 superconformal invariance.^{14,15} This method is manifestly spacetime supersymmetric like the light-cone Green-Schwarz method, but requires no operator insertions at the interaction points. Although the scattering amplitudes are calculated by evaluating correlation functions on N=2 super-Riemann surfaces, there is no ambiguity coming from the choice of anti-commuting moduli since the dilaton tadpole diagram vanishes by spacetime supersymmetry,[†] and there is therefore no need to introduce a cutoff in the moduli space. Using standard BRST techniques, Green-Schwarz superstring scattering amplitudes with an arbitrary number of loops and external states can be explicitly evaluated. These amplitudes agree with those obtained using the manifestly unitary light-cone Green-Schwarz formalism if one makes a simple conjecture concerning the contribution of the light-cone gauge contact terms (although a similar conjecture in the NSR formalism can be proven to be true, a proof in the Green-Schwarz formalism is still lacking).

The only disadvantage of this new method is that in order to express the N=2 twistor-string action in terms of free fields, the manifest SO(9,1) super-Poincaré invariance has to be broken down to an SU(4)xU(1) subgroup. However, it will be shown in this paper that the twistor-string free fields can be combined into N=2 chiral and anti-chiral superfields, $X_{\pm}^{\mu}$ and $\Psi_{\pm}^{\alpha}$, that transform covariantly as SO(9,1) vectors and spinors under commutation with the SO(9,1) Lorentz generators, $M^{\mu\nu} = \int dz d^2\kappa X_{+}^{\mu} X_{-}^{\nu}$. These covariantly transforming superfields do not have free-field operator product expansions, but instead obey the relation, $\Psi_{-}^{\alpha} \Psi_{+}^{\beta} = \gamma_{\mu}^{\alpha\beta} (X_{+}^{\mu} - X_{-}^{\mu})$, which is a ten-dimensional generalization of the usual analyticity condition in four dimensions.^{17,18}

The gauge-fixed physical vertex operators for the massless states that were introduced in reference 15 can now easily be extended to manifestly Lorentz-covariant vertex operators. The right-moving part

[†] The spacetime supersymmetry generators in the twistor-string formalism contain no ghost contributions, so the dilaton tadpole diagram can be shown to vanish by writing the zero-momentum dilaton vertex operator as the contour integral of a spacetime supersymmetry generator around a zero-momentum dilatino vertex operator, and pulling the contour off the back of the surface. In the Neveu-Schwarz-Ramond formalism, the ghost contributions to the spacetime supersymmetry generators have unwanted poles,² so the same argument¹⁶ implies only that the dilaton tadpole diagram is a total derivative in moduli space.⁴

of the bosonic vertex operator is simply $\eta_\mu \int dz d^2 \kappa X_+^\mu \exp(ik \cdot X_-)$, while the right-moving part of the fermionic vertex operator is $u_\alpha \int dz d^2 \kappa \Psi_+^\alpha \exp(ik \cdot X_-)$. Because the $SO(9,1)$ super-Poincaré generators are superconformally invariant, the scattering amplitudes obtained by evaluating correlation functions of these covariant vertex operators on $N=2$ super-Riemann surfaces are guaranteed to be super-Poincaré covariant.

In order to explicitly evaluate these correlation functions, the $X_\pm^\mu$ and $\Psi_\pm^\alpha$ superfields must be re-expressed in terms of the original free fields, whose correlation functions on arbitrary genus surfaces were calculated in reference 15. Although this procedure breaks the manifest $SO(9,1)$ invariance down to an $SU(4) \times U(1)$ subgroup, it is straightforward to use knowledge of the underlying larger invariance to write the resulting scattering amplitudes in manifestly Lorentz-covariant notation.

In the conclusion of this paper, the issue of finiteness will be briefly discussed, as well as the possibility of generalizing the amplitude calculations to non-flat target-space backgrounds.

II. Free Fields of the $N=2$ Twistor-String

The free-field action for the $N=2$ twistor-string corresponding to the heterotic Green-Schwarz superstring is:¹⁴

$$S = \int dz d\bar{z} d\kappa^+ d\kappa^- \left[\frac{1}{2} (X^{+\bar{l}} \partial_{\bar{z}} X^{-l} - X^{-l} \partial_{\bar{z}} X^{+\bar{l}}) + W^- \partial_{\bar{z}} \Psi^+ - W^+ \partial_{\bar{z}} \Psi^- + \Phi^{+q} \Phi^{-q} \right] \quad (II.1)$$

where $[z, \kappa^+, \kappa^-]$ and $[\bar{z}]$ are holomorphic and anti-holomorphic coordinates on the $N=(2,0)$ super-Riemann surface (although this Euclidean surface is not well-defined since $(\kappa^\pm)^*$ does not exist, independence of the right and left-moving sectors allows heterotic superstring amplitudes to be defined in the usual way by taking the holomorphic square-root of non-heterotic amplitudes), $D_\pm = \partial_{\kappa^\pm} + \frac{1}{2} \kappa^\mp \partial_z$, $X^{+\bar{l}}$ and X^{-l} for $l = 1$ to 4 describe the transverse degrees of freedom and transform as $\bar{4}_{+\frac{1}{2}}$ and $4_{-\frac{1}{2}}$ under the remaining $SU(4) \times U(1)$ invariance, $D_\pm W^\pm$ and $\Psi^\pm$ describe the longitudinal degrees of freedom and transform as $1_{\pm\frac{1}{2}}$ under $SU(4) \times U(1)$, and $\Phi^{\pm q}|_{\kappa^+ = \kappa^- = 0}$ for $q=1$ to 16 describe the heterotic lattice and are $SU(4) \times U(1)$ scalars. These $N=(2,0)$ superfields are restricted by the chirality constraints:

$$D_- X^{+\bar{l}} = D_+ X^{-l} = D_- \Psi^+ = D_+ \Psi^- = D_- \Phi^{+q} = D_+ \Phi^{-q}; \quad (II.2)$$

the global constraint:

$$D_+ \Psi^+ D_- \Psi^- - \Psi^+ \partial_z \Psi^- - \Psi^- \partial_z \Psi^+ = \partial_z X^{9+0} \text{ for some real superfield } X^{9+0}; \quad (II.3)$$

and the super-Virasoro constraints:

$$D_+ W^+ D_- \Psi^- - D_- W^- D_+ \Psi^+ + D_+ X^{+\bar{l}} D_- X^{-l} = \kappa^+ \kappa^- [\partial_{\bar{z}} X^{+\bar{l}} \partial_{\bar{z}} X^{-l} + \partial_{\bar{z}} X^{9-0} \partial_{\bar{z}} X^{9+0} + \frac{1}{2} (\Phi^{+q} \partial_{\bar{z}} \Phi^{-q} + \Phi^{-q} \partial_{\bar{z}} \Phi^{+q})] = 0, \quad (II.4)$$

where $X^{9-0} \equiv \frac{D_+ W^+}{D_+ \Psi^+} + \frac{D_- W^-}{D_- \Psi^-}$.

This free-field action is obtained by gauge-fixing the manifestly super-Poincaré invariant N=(2,0) twistor-string action, which has been shown to have the same classical degrees of freedom as the heterotic Green-Schwarz superstring action.¹⁹ Evidence for the quantum consistency of the free-field action comes from the vanishing of the superconformal anomaly and from the fact that after integrating out the $\Psi^\pm$ and $W^\pm$ superfields, one recovers the usual light-cone gauge Green-Schwarz action, including the interaction-point operators.¹⁴

In order to eliminate unnecessary notation, only the heterotic superstring will be discussed in this paper although all techniques straightforwardly generalize to the non-heterotic cases. For example, the free-field action for the non-heterotic superstring is obtained from equation (II.1) by extending the worldsheet to an N=(2,2) surface and replacing the left-moving superfields, $\Phi^{\pm q}$, with $\bar{W}^\pm$ and $\bar{\Psi}^\pm$ superfields.¹⁵ The only subtle point is that in the Type IIA theory, the roles of X^{9+0} and X^{9-0} are reversed in the two different sectors, i.e. the right-moving global constraint defines X^{9+0} , whereas the left-moving global constraint defines X^{9-0} .

As discussed in reference 15, it is convenient to solve the global constraint of equation (II.3) by bosonizing the components of $\Psi^\pm = \psi^\pm + \kappa^\pm \lambda^\pm$ and $D_\pm W^\pm = w^\pm + \kappa^\mp \varepsilon^\pm$ in the following way:

$$\lambda^+ = (\partial_z x^{9+0} + \frac{1}{2} \psi^+ \partial_z \psi^- + \frac{1}{2} \psi^- \partial_z \psi^+) e^{h^+} + e^{-h^-}, \quad \lambda^- = e^{-h^+} \quad (II.5)$$

$$w^+ = e^{h^+} (\partial_z h^+ + \partial_z h^- + x^{9-0} (\partial_z x^{9+0} + \frac{1}{2} \psi^+ \partial_z \psi^- + \frac{1}{2} \psi^- \partial_z \psi^+)) + x^{9-0} e^{-h^-}, \quad w^- = x^{9-0} e^{-h^+},$$

where $h^\pm$ are chiral bosons with screening charge +1 that satisfy $h^+(y)h^-(z) \rightarrow \log(y-z)$ as $y \rightarrow z$.

In terms of these bosonized fields, the right-handed super-Virasoro constraints of equation (II.4) take the form:

$$\begin{aligned} T &= \Gamma^{+\bar{l}} \Gamma^{-l} - \partial_z h^+ + \partial_z h^-, \quad G_- = \partial_z x^{+\bar{l}} \Gamma^{-l} + (\hat{\varepsilon}^+ + \frac{1}{2} \psi^+ \partial_z x^{9-0}) e^{-h^+}, \quad G_+ = \partial_z x^{-l} \Gamma^{+\bar{l}} + \\ &(\hat{\varepsilon}^- + \frac{1}{2} \partial_z x^{9-0} \psi^-) ((\partial_z x^{9+0} + \frac{1}{2} \psi^+ \partial_z \psi^- + \frac{1}{2} \psi^- \partial_z \psi^+) e^{h^+} + e^{-h^-}) - e^{h^+} ((\partial_z h^+ + \partial_z h^-) \partial_z \psi^- + \frac{3}{4} \partial_z^2 \psi^-), \\ L &= \partial_z x^{+\bar{l}} \partial_z x^{-l} - \frac{1}{2} (\Gamma^{+\bar{l}} \partial_z \Gamma^{-l} + \Gamma^{-l} \partial_z \Gamma^{+\bar{l}}) - \hat{\varepsilon}^+ \partial_z \psi^- - \hat{\varepsilon}^- \partial_z \psi^+ + \partial_z h^+ \partial_z h^- + \frac{1}{2} (\partial_z^2 h^+ + \partial_z^2 h^-), \end{aligned} \quad (II.6)$$

where $X^{+\bar{l}} = x^{+\bar{l}} + \kappa^+ \Gamma^{+\bar{l}}$, $X^{-l} = x^{-l} + \kappa^- \Gamma^{-l}$, and $\hat{\varepsilon}^\pm \equiv \varepsilon^\pm - \frac{1}{2} \partial_z x^{9-0} \psi^\pm - x^{9-0} \partial_z \psi^\pm$.

III. SO(9,1) Lorentz Generators

The simplest way to construct the SO(9,1) Lorentz generators out of these free fields is to look for a generalization of the operator, $m^{\mu\nu} = \int dz \, x^\mu \partial_z x^\nu$, that is N=(2,0) superconformally invariant (the left-moving contribution to the Lorentz generator, $\int d\bar{z} \, x^\mu \partial_{\bar{z}} x^\nu$, is already superconformally invariant). Since for the 15 SU(4) generators that preserve the gauge-fixing, these generalized operators are simply

$M^{+\bar{l}-m} = \int dz d^2\kappa X^{+\bar{l}} X^{-m}$, it is natural to look for chiral and anti-chiral superfields, $X_{\pm}^{\mu}$, that contain x^{μ} in their $\kappa^{\pm} = 0$ component.

Using the free-field commutation relations, it is easy to show that $[G_+, x_+^{\mu}] = [G_-, x_-^{\mu}] = 0$, where

$$\begin{aligned} x_+^{9+0} &\equiv x^{9+0} - \frac{1}{2}\psi^+\psi^-, \quad x_+^{9-0} \equiv x^{9-0} + e^{h^++h^-}, \quad x_+^{+\bar{l}} \equiv x^{+\bar{l}} - e^{h^-}\psi^+\Gamma^{+\bar{l}}, \quad x_+^{-l} \equiv x^{-l}; \\ x_-^{9+0} &\equiv x^{9+0} + \frac{1}{2}\psi^+\psi^-, \quad x_-^{9-0} \equiv x^{9-0}, \quad x_-^{+\bar{l}} \equiv x^{+\bar{l}}, \quad x_-^{-l} \equiv x^{-l} - e^{h^+}\psi^-\Gamma^{-l}; \end{aligned} \quad (III.1)$$

and $G_{\pm}$ are defined in equation (II.6). Therefore, the desired chiral and anti-chiral superfields are

$$X_+^{\mu}(z^-, \kappa^-) = x_+^{\mu}(z^-) + \kappa^- [G_-, x_+^{\mu}(z^-)] \quad \text{and} \quad X_-^{\mu}(z^+, \kappa^+) = x_-^{\mu}(z^+) + \kappa^+ [G_+, x_-^{\mu}(z^+)], \quad (III.2)$$

where $z^{\pm} \equiv z \pm \frac{1}{2}\kappa^+\kappa^-$. Note that $x_+^{\mu} = [G_+, x_-^{\mu}\psi^+e^{h^-}]$, $x_-^{\mu} = [G_-, x_+^{\mu}\psi^-e^{h^+}]$, and $X_+^{\mu}(y^-, \xi^-) X_-^{\nu}(z^+, \kappa^+) \rightarrow \eta^{\mu\nu} \log(y^- - z^+ - \xi^- \kappa^+)$ as $y \rightarrow z$.

It can now be straightforwardly checked using equations (II.6), (III.1), and (III.2), that the 45 operators,

$$M^{\mu\nu} = \int dz d^2\kappa X_+^{\mu} X_-^{\nu} \quad (= - \int dz d^2\kappa X_+^{\nu} X_-^{\mu} \text{ for } \mu \neq \nu), \quad (III.3)$$

are superconformally invariant, commute with each other to form an $\text{SO}(9,1)$ algebra, and transform X_+^{μ} and X_-^{μ} like covariant $\text{SO}(9,1)$ vectors.

By commuting $M^{\mu\nu}$ with the chiral and anti-chiral superfields $\Psi^{\pm}$, one finds that Ψ^+ transforms as the 1_{+1} component of an $\text{SO}(9,1)$ Weyl spinor superfield Ψ_-^{α} , and Ψ^- transforms as the 1_{-1} component of an $\text{SO}(9,1)$ Weyl spinor superfield Ψ_+^{α} , where

$$\begin{aligned} \Psi_-^{\alpha}(z^+, \kappa^+) &= \psi_-^{\alpha}(z^+) + \kappa^+ \{G_+, \psi_-^{\alpha}(z^+)\}, \quad \Psi_+^{\alpha}(z^-, \kappa^-) = \psi_+^{\alpha}(z^-) + \kappa^- \{G_-, \psi_+^{\alpha}(z^-)\}; \quad (III.4) \\ \psi_-^+ &= \psi^+, \quad \psi_-^{lm} = e^{h^+-h^-} \psi^- \Gamma^{-l} \Gamma^{-m}, \quad \psi_-^- = \frac{1}{24} e^{2h^+-2h^-} \psi^- \partial_z \psi^- \hat{\varepsilon}^- \epsilon_{lmnp} \Gamma^{-l} \Gamma^{-m} \Gamma^{-n} \Gamma^{-p}, \\ \psi_-^{+l} &= e^{h^+} \Gamma^{-l}, \quad \psi_-^{-\bar{l}} = \frac{1}{6} e^{2h^+-h^-} \psi^- \hat{\varepsilon}^- \epsilon_{lmnp} \Gamma^{-m} \Gamma^{-n} \Gamma^{-p}; \\ \psi_+^+ &= \frac{1}{24} e^{2h^--2h^+} \psi^+ \partial_z \psi^+ \hat{\varepsilon}^+ \epsilon^{lmnp} \Gamma^{+l} \Gamma^{+m} \Gamma^{+n} \Gamma^{+p}, \quad \psi_+^{lm} = \frac{1}{2} e^{h^--h^+} \psi^+ \epsilon^{lmnp} \Gamma^{+n} \Gamma^{+p}, \quad \psi_+^- = \psi^-, \\ \psi_+^{+l} &= \frac{1}{6} e^{2h^--h^+} \psi^+ \hat{\varepsilon}^+ \epsilon^{lmnp} \Gamma^{+m} \Gamma^{+n} \Gamma^{+p}, \quad \psi_+^{-\bar{l}} = e^{h^-} \Gamma^{+\bar{l}}; \end{aligned}$$

and the sixteen components of the $\text{SO}(9,1)$ Weyl spinor have been split up into $[1_{+1}, 6_0, 1_{-1}, 4_{+1}, \bar{4}_{-1}]$ representations of $\text{SU}(4) \times \text{U}(1)$.

As was mentioned in the introduction, these spinor superfields obey the identity,

$$\Psi_-^{\alpha} \Psi_+^{\beta} = (X_+^{\mu} - X_-^{\mu}) \gamma_{\mu}^{\alpha\beta} \quad (III.5),$$

which can easily be checked using their free-field expansions. This identity is an obvious generalization of the four-dimensional analyticity condition,^{17,18} $\theta^{\alpha} \bar{\theta}^{\dot{\alpha}} = (x_+^{\mu} - x_-^{\mu}) \gamma_{\mu}^{\alpha\dot{\alpha}}$, however because the ten-dimensional identity contains 256 components, it can not be solved using purely classical fields.

IV. Covariant Vertex Operators

Using the $X_{\pm}^{\mu}$ and $\Psi_{\pm}^{\alpha}$ superfields, it is easy to construct the superconformally invariant vertex operators that covariantly describe the massless bosonic and fermionic states. The massless bosonic vertex operator is

$$\eta_{\mu} \int d^2 z d^2 \kappa X_{+}^{\mu} e^{ik \cdot X} V_L \quad (= -\eta_{\mu} \int d^2 z d^2 \kappa X_{-}^{\mu} e^{ik \cdot X} V_L), \quad (IV.1)$$

where the left-moving contribution to the vertex operator, V_L , is constructed in the usual way for the heterotic superstring²⁰ out of $\partial_{\bar{z}} X_{+}^{\mu}$ and $\Phi^{\pm q}$ (note that only the $\kappa^{\pm} = 0$ components of these superfields contribute), and $k^2 = \eta \cdot k = 0$. This vertex operator is invariant under the gauge transformations, $\eta^{\mu} \rightarrow \eta^{\mu} + \Lambda k^{\mu}$, and after fixing the gauge $\eta^{9+0} = 0$ and replacing the $\int d^2 z$ integration with $c\bar{c}$ ghosts, it coincides with the massless bosonic vertex operator of reference 15 in the picture with ghost-number (1,1) and instanton-number zero (the instanton-number of a vertex operator is defined as its eigenvalue under commutation with the charge, $K = \frac{1}{2} \int dz (\hat{\varepsilon}^{+} \psi^{-} - \hat{\varepsilon}^{-} \psi^{+} + \partial_z h^{-} - \partial_z h^{+})$, and can be shifted by an arbitrary integer by attaching instanton-number-changing operators to the vertex operator).

Two covariant choices for the massless fermionic vertex operator are

$$u_{\alpha} \int d^2 z d^2 \kappa \Psi_{-}^{\alpha} e^{ik \cdot X} V_L \quad \text{or} \quad u_{\alpha} \int d^2 z d^2 \kappa \Psi_{+}^{\alpha} e^{ik \cdot X} V_L, \quad (IV.2)$$

where $k^2 = 0$. These vertex operators are invariant under the gauge transformations, $u_{\alpha} \rightarrow u_{\alpha} + k_{\mu} \Lambda^{\beta} \gamma_{\alpha\beta}^{\mu}$, and after fixing the gauge $(\gamma^{9+0} u)^{\alpha} = 0$ and replacing the $\int d^2 z$ integration with $c\bar{c}$ ghosts, they coincide with the massless fermionic vertex operator of reference 15, either in the picture with ghost-number (1,1) and instanton-number $-\frac{1}{2}$, or in the picture with ghost-number (1,1) and instanton-number $+\frac{1}{2}$.

Since the usual fermionic field, ζ^{α} , satisfies $k_{\mu} \zeta^{\alpha} \gamma_{\alpha\beta}^{\mu} = 0$ and has no gauge invariances, it is clear that ζ^{α} corresponds to $k^{\mu} u_{\beta} \gamma_{\mu}^{\alpha\beta}$. One therefore still needs to construct the vertex operator for the fermionic field when $k^{\mu} = 0$. Two choices for this zero-momentum vertex operator are

$$\zeta^{\alpha} \int d^2 z d^2 \kappa \gamma_{\mu, \alpha\beta} X_{+}^{\mu} \Psi_{-}^{\beta} V_L \quad \text{or} \quad \zeta^{\alpha} \int d^2 z d^2 \kappa \gamma_{\mu, \alpha\beta} X_{-}^{\mu} \Psi_{+}^{\beta} V_L, \quad (IV.3)$$

out of which can be extracted two choices for the covariant spacetime supersymmetry generators,

$$S_{-, \alpha} = \frac{2}{5} \int dz d^2 \kappa \gamma_{\mu, \alpha\beta} X_{+}^{\mu} \Psi_{-}^{\beta} \quad \text{or} \quad S_{+, \alpha} = \frac{2}{5} \int dz d^2 \kappa \gamma_{\mu, \alpha\beta} X_{-}^{\mu} \Psi_{+}^{\beta}, \quad (IV.4)$$

which satisfy $\{S_{-, \alpha}, S_{+, \beta}\} = 2 \int dz \partial_z x_{\mu} \gamma_{\alpha\beta}^{\mu}$. It is easily verified that commutation with $S_{\pm, \alpha}$ takes the fermionic vertex operator of instanton-number $\mp \frac{1}{2}$ and spinor polarization u_{β} into the bosonic vertex operator of vector polarization $\eta^{\mu} = (\gamma^{\mu} \gamma^{\nu} u)_{\alpha} k_{\nu}$, and takes the bosonic vertex operator of vector polarization η^{μ} into the fermionic vertex operator of instanton-number $\pm \frac{1}{2}$ and spinor-polarization $u_{\beta} = \gamma_{\alpha\beta}^{\mu} \eta_{\mu}$.

It is interesting to note that the integrands of the spacetime supersymmetry generators,

$$W_{-, \alpha} = \gamma_{\mu, \alpha\beta} X_{+}^{\mu} \Psi_{-}^{\beta} \quad \text{and} \quad W_{+, \alpha} = \gamma_{\mu, \alpha\beta} X_{-}^{\mu} \Psi_{+}^{\beta}, \quad (IV.5)$$

satisfy the free-field behavior

$$D_+ W_-(y^-, \xi^-) \Psi_+(z^-, \kappa^-) \rightarrow (\xi^- - \kappa^-)/(y^- - z^-) \text{ and } D_- W_+(y^+, \xi^+) \Psi_-(z^+, \kappa^+) \rightarrow (\xi^+ - \kappa^+)/(y^+ - z^+)$$

as $y \rightarrow z$, and therefore provide an $N=(2,0)$ generalization of the twistor equation,^{21,22} $w_\alpha = \gamma_{\mu,\alpha\beta} x^\mu \lambda^\beta$.

V. Lorentz-Covariant Scattering Amplitudes

Scattering amplitudes can now be calculated by evaluating correlation functions of these covariant vertex operators on $N=2$ super-Riemann surfaces and integrating over the global super-moduli of the surfaces. By choosing half of the fermionic vertex operators to have instanton-number $+\frac{1}{2}$ and the other half to have instanton-number $-\frac{1}{2}$, the $N=2$ super-Riemann surfaces can be restricted to those with vanishing instanton-number (in order to get a non-zero correlation function, the instanton-number of the surface must equal the sum of the instanton-numbers of the vertex operators). The choice of which fermionic vertex operators have $+\frac{1}{2}$ or $-\frac{1}{2}$ instanton-number does not affect the scattering amplitude since different choices correspond to different locations for the picture-changing and instanton-number-changing operators (see reference 15 for more details). Similarly, the choice of using $S_{+,\alpha}$ or $S_{-,\alpha}$ does not affect the scattering amplitude as long as the instanton-number of the surface is set equal to the total instanton-number of the vertex operators plus the total instanton-number of the spacetime supersymmetry generators.

Since $M^{\mu\nu}$ and $S_{\pm,\alpha}$ of equations (III.3) and (IV.4) are superconformally invariant and covariantly transform the vertex operators, these scattering amplitudes are proven to be $SO(9,1)$ super-Poincaré covariant.

At the present time, the only known way to evaluate correlation functions of the covariant vertex operators is to re-express the vertex operators in terms of the original set of free fields, and to use the free-field correlation functions that were evaluated on arbitrary genus surfaces in reference 15. Although this procedure breaks the manifest Lorentz covariance down to an $SU(4) \times U(1)$ subgroup, knowledge of the underlying larger invariance can be used to express the resulting scattering amplitudes in manifestly Lorentz-covariant notation.

The first step in covariantizing the calculation is to introduce an $SO(9,1)$ pure spinor²³ and its complex conjugate, v^α and $\bar{v}^\alpha$, and a real $SO(9,1)$ light-like vector, m^μ , which satisfy

$$v^\alpha \gamma_{\alpha\beta}^\mu v^\beta = \bar{v}^\alpha \gamma_{\alpha\beta}^\mu \bar{v}^\beta = m^\mu m_\mu = 0 \quad \text{and} \quad v^\alpha \gamma_{\alpha\beta}^\mu \bar{v}^\beta m_\mu = 1. \quad (V.1)$$

With the help of these projection operators, the free fields $\Gamma^{+\bar{l}}$ and Γ^{-l} can be treated as four complex components of $SO(9,1)$ vectors Γ_-^μ and Γ_+^μ (all other components of the vectors will be projected out), while the free fields $\psi^\pm$, $\bar{\varepsilon}^\pm$, $h^\pm$, as well as all ghost fields (see reference 15 for details on the ghost fields), can be

treated as SO(9,1) scalars. For example, the $\kappa^\pm = 0$ component of the covariant superfields X_-^μ and Ψ_-^α can be written as

$$\begin{aligned} x_-^\mu &= x^\mu + \frac{1}{2}\psi^+\psi^-(v\gamma^\mu\bar{v}) - e^{h^+}\psi^-m^\nu\Gamma_+^\rho(v\gamma^\mu\gamma_\nu\gamma_\rho\bar{v}), \\ \psi_-^\alpha &= \psi^+\bar{v}^\alpha + e^{h^+}m^\mu\Gamma_+^\nu(\gamma_\mu\gamma_\nu\bar{v})^\alpha + \frac{1}{4}e^{h^+-h^-}\psi^-m^\rho\Gamma_+^\sigma\Gamma_+^\tau(\bar{v}\gamma_\mu\gamma_\nu\gamma_\rho\gamma_\sigma\gamma_\tau\bar{v})(\gamma^\mu\gamma^\nu v)^\alpha + \\ &\quad \frac{1}{6}e^{2h^+-h^-}\psi^-\hat{\varepsilon}^-m^\mu\Gamma_+^\nu\Gamma_+^\rho\Gamma_+^\sigma(\gamma_\mu\gamma_\nu\gamma_\rho\gamma_\sigma\bar{v})^\alpha + \frac{1}{24}e^{2h^+-2h^-}\psi^-\partial_z\psi^-\hat{\varepsilon}^-m^\mu\Gamma_+^\nu\Gamma_+^\rho\Gamma_+^\sigma\Gamma_+^\tau(\bar{v}\gamma_\mu\gamma_\nu\gamma_\rho\gamma_\sigma\gamma_\tau\bar{v})v^\alpha, \end{aligned} \quad (V.2)$$

and the term $\partial_z x^{-l}\Gamma^{+\bar{l}}$ in G_+ of equation (III.6) can be written as $\partial_z x^\mu m^\nu\Gamma_-^\rho(\bar{v}\gamma_\mu\gamma_\nu\gamma_\rho v)$.

After evaluating correlation functions of the free fields as in reference 15, one is left with an expression for the scattering amplitude that is manifestly Lorentz covariant, but which is a polynomial in the three projection operators, v^α , $\bar{v}^\alpha$, and m^μ . However, the knowledge that the scattering amplitude is Lorentz covariant even without transforming the projection operators implies that the projection operators only occur in Lorentz-invariant combinations. Therefore, replacing all monomials of projection operators by their Lorentz-invariant component does not affect the scattering amplitude, since all non-Lorentz-invariant components must cancel out. So covariantization of the Green-Schwarz superstring amplitudes has been reduced to the problem of extracting the Lorentz-invariant component of the monomial $v^{\alpha_1}\dots v^{\alpha_L}\bar{v}^{\beta_1}\dots\bar{v}^{\beta_M}m^{\mu_1}\dots m^{\mu_N}$, where v^α , $\bar{v}^\alpha$, and m^μ satisfy equation (V.1). Note that this covariantization procedure requires that the amplitude does not contain negative powers of the projection operators, and therefore is not useful in the light-cone or semi-light-cone Green-Schwarz formalisms where $(k^{9+0})^{-1}$ factors are present.

Since under a U(1) rotation, $v^\alpha \rightarrow e^{i\phi}v^\alpha$, $\bar{v}^\alpha \rightarrow e^{-i\phi}\bar{v}^\alpha$, $m^\mu \rightarrow m^\mu$; while under an SO(1,1) boost, $v^\alpha \rightarrow e^\phi v^\alpha$, $\bar{v}^\alpha \rightarrow e^\phi \bar{v}^\alpha$, $m^\mu \rightarrow e^{-2\phi}m^\mu$; the Lorentz-invariant component of a monomial is zero unless there are an equal number of v^α 's, $\bar{v}^\alpha$'s and m^μ 's. The Lorentz-invariant component, $P_N^{\alpha_1\dots\alpha_N\beta_1\dots\beta_N\mu_1\dots\mu_N}$, of all such contributing monomials, $v^{\alpha_1}\dots v^{\alpha_N}\bar{v}^{\beta_1}\dots\bar{v}^{\beta_N}m^{\mu_1}\dots m^{\mu_N}$, can be obtained from the formula:

$$P_N^{\alpha_1\dots\alpha_N\beta_1\dots\beta_N\mu_1\dots\mu_N} = c_N \sum_{\text{symmetrized}} \quad (V.3)$$

$$[(N!)^{-1}\gamma^{\mu_1,\alpha_1\beta_1}\dots\gamma^{\mu_N,\alpha_N\beta_N} + \frac{(2(N-2)!)^{-3}}{N+2}\gamma_\nu^{\alpha_1\alpha_2}\gamma_\rho^{\beta_1\beta_2}(\eta^{\nu\rho}\eta^{\mu_1\mu_2} - 2\eta^{\nu\mu_1}\eta^{\rho\mu_2})P_{N-2}^{\alpha_3\dots\alpha_N\beta_3\dots\beta_N\mu_3\dots\mu_N}],$$

where $\sum$ means to symmetrize independently in the α , β , and μ indices to give $(N!)^3$ terms, $P_0 = 1$, $P_1^{\alpha\beta\mu} = \frac{1}{160}\gamma^{\mu,\alpha\beta}$, and c_N is fixed by the requirement that $\gamma_{\mu_1,\alpha_1\beta_1}\dots\gamma_{\mu_N,\alpha_N\beta_N}P_N^{\alpha_1\dots\alpha_N\beta_1\dots\beta_N\mu_1\dots\mu_N} = 1$. It is straightforward to check that $P_N^{\alpha_1\dots\alpha_N\beta_1\dots\beta_N\mu_1\dots\mu_N}$ is the unique Lorentz-invariant tensor which is symmetric in the α , β , and μ indices, which vanishes when contracted with $\gamma_{\alpha_1\alpha_2}^\sigma$, $\gamma_{\beta_1\beta_2}^\sigma$, or $\eta_{\mu_1\mu_2}$, and which is normalized to one when all indices are contracted with γ -matrices. So after replacing all contributing monomials by P_N in the expression for the scattering amplitude that is calculated using the free-field description of the covariant vertex operators, the Green-Schwarz superstring amplitudes are written in manifestly Lorentz-covariant notation.

VI. Conclusion

After constructing $SO(9,1)$ super-Poincaré generators and manifestly covariant vertex operators, it was proved in this paper that the earlier calculated multiloop Green-Schwarz superstring scattering amplitudes are super-Poincaré invariant, and it was shown how to write these amplitudes in manifestly Lorentz-covariant form.

One possible use for these amplitudes is to analyze their finiteness properties. Using the Neveu-Schwarz-Ramond method, it is difficult to show that the dilaton tadpole diagram vanishes,⁵ while using the light-cone Green-Schwarz method, it is difficult to show that the contact terms precisely cancel the divergences when two or more interaction points coalesce.⁹ Since neither of these difficulties are present using the twistor-string method, it should be possible to check for finiteness by explicitly looking for divergences in the scattering amplitude expressions.

In the construction of the super-Poincaré generators and covariant vertex operators, the $N=2$ chiral and anti-chiral superfields $X_{\pm}^{\mu}$ and $\Psi_{\pm}^{\alpha}$ played a prominent role. It is tempting to suggest that in a non-flat target-space background, these $N=2$ superfields provide the phase space for the ten-dimensional supergravity and super-Yang-Mills fields.²⁴ In flat space, the superfields obey the relations, $\Psi_{-}^{\alpha}\Psi_{+}^{\beta} = (X_{+}^{\mu} - X_{-}^{\mu})\gamma_{\mu}^{\alpha\beta}$ from equation (III.5), and $W_{+,\alpha} = \gamma_{\mu,\alpha\beta}X_{-}^{\mu}\Psi_{+}^{\beta}$ from equation (IV.5), where $X_{+}^{\mu}(y^{-}, \xi^{-})X_{-}^{\nu}(z^{+}, \kappa^{+}) \rightarrow \log(y^{-} - z^{+} - \xi^{-}\kappa^{+})$ and $D_{-}W_{+}(y^{+}, \xi^{+})\Psi_{-}(z^{+}, \kappa^{+}) \rightarrow (\xi^{+} - \kappa^{+})/(y^{+} - z^{+})$. Perhaps in a curved target-space background, these superfield relations are modified in the same way that the four-dimensional analyticity condition, $\theta^{\alpha}\bar{\theta}^{\dot{\alpha}} = (x_{+}^{\mu} - x_{-}^{\mu})\gamma_{\mu}^{\alpha\dot{\alpha}}$, is modified to $H(x, \theta, \bar{\theta})^{\alpha\dot{\alpha}} = (x_{+}^{\mu} - x_{-}^{\mu})\gamma_{\mu}^{\alpha\dot{\alpha}}$ in the presence of supergravity.²⁵

Acknowledgements

I would like to thank M. Green, P. Howe, S. Mandelstam, W. Siegel, J. Taylor, and P. Townsend for useful discussions, and the SERC for its financial support.

References

- (1) Friedan, D., Martinec, E., and Shenker, S., Nucl.Phys.B271 (1986), p.93.
- (2) Verlinde, E. and Verlinde, H., Phys.Lett.B192 (1987), p.95.
- (3) Aoki, K., D'Hoker, E., and Phong, D.H., Nucl.Phys.B342 (1990), p.149.
- (4) Atick, J. and Sen, A., Nucl.Phys.B296 (1988), p.157.
- (5) Mandelstam, S., "The n-loop Amplitude: Explicit Formulas, Finiteness, and Absence of Ambiguities", preprint UCB-PTH-91/53, October 1991.
- (6) Iengo, R. and Zhu, C.-J., Phys.Lett.B212 (1988), p.313.
- (7) Green, M.B. and Schwarz, J.H., Nucl.Phys.B243 (1984), p.475.

- (8) Mandelstam,S., Prog.Theor.Phys.Suppl.86 (1986), p.163.
- (9) Restuccia,A. and Taylor,J.G., Phys.Rep.174 (1989), p.283.
- (10) Greensite,J. and Klinkhamer,F.R., Nucl.Phys.B291 (1987), p.557.
- (11) Mandelstam,S., private communication.
- (12) Carlip,S., Nucl.Phys.B284 (1987), p.365.
- (13) Gilbert,G. and Johnston,D., Phys.Lett.B205 (1988), p.273.
- (14) Berkovits,N., Nucl.Phys.B379 (1992), p.96., hep-th bulletin board 9201004.
- (15) Berkovits,N., “Calculation of Green-Schwarz Superstring Amplitudes using the N=2 Twistor-String Formalism”, SUNY at Stonybrook preprint ITP-SB-92-42, August 1992, hep-th bulletin board 9208035.
- (16) Martinec,E., Phys.Lett.B171 (1986), p.189.
- (17) Delduc,F. and Sokatchev,E., Class.Quant.Grav. 9 (1992), p.361.
- (18) Ivanov,E.A. and Kapustnikov,A.A., Phys.Lett.B267 (1991), p.175.
- (19) Tonin,M., Phys.Lett.B266 (1991), p.312.
- (20) Gross,D.J., Harvey,J.A., Martinec,E., and Rohm,R., Nucl.Phys.B256 (1985), p.253.
- (21) Penrose,R. and MacCallum,M.A.H., Phys.Rep.6C (1972), p.241.
- (22) Sorokin,D.P., Tkach,V.I., Volkov,D.V., and Zheltukhin,A.A., Phys.Lett.B216 (1989), p.302.
- (23) Howe,P.S., Phys.Lett.B258 (1991), p.141.
- (24) Gates,S.J.Jr. and Nishino,H., Class.Quant.Grav. 3 (1986), p.39.
- (25) Ogievetsky,V.I. and Sokatchev,E.S., Phys.Lett.79 (1978), p.222.